



ACCESS & REIMBURSEMENT

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.

Table of Contents

GIVLAARI® (givosiran) Indication and Important Safety Information	3
Alnylam Assist™ Services Overview	4
How to Complete the GIVLAARI® (givosiran) Start Form	11
Overview of Acquisition Process for GIVLAARI® (givosiran)	18
Billing and Coding Overview	20
Billing and Coding—Physician Office	22
Billing and Coding—Hospital Outpatient Department	27
Alnylam Assist™ Copay Claim Submission Guide	32
Dosing and Administration Guide	35
Provider Readiness Guide	41

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





INDICATION

GIVLAARI® (givosiran) is indicated for the treatment of adults with acute hepatic porphyria (AHP).

IMPORTANT SAFETY INFORMATION

Contraindications

GIVLAARI is contraindicated in patients with known severe hypersensitivity to givosiran. Reactions have included anaphylaxis.

Anaphylactic Reaction

Anaphylaxis has occurred with GIVLAARI treatment (<1% of patients in clinical trials). Ensure that medical support is available to appropriately manage anaphylactic reactions when administering GIVLAARI. Monitor for signs and symptoms of anaphylaxis. If anaphylaxis occurs, immediately discontinue administration of GIVLAARI and institute appropriate medical treatment.

Hepatic Toxicity

Transaminase elevations (ALT) of at least 3 times the upper limit of normal (ULN) were observed in 15% of patients receiving GIVLAARI in the placebo-controlled trial. Transaminase elevations primarily occurred between 3 to 5 months following initiation of treatment.

Measure liver function tests prior to initiating treatment with GIVLAARI, repeat every month during the first 6 months of treatment, and as clinically indicated thereafter. Interrupt or discontinue treatment with GIVLAARI for severe or clinically significant transaminase elevations. In patients who have dose interruption and subsequent improvement, reduce the dose to 1.25 mg/kg once monthly. The dose may be increased to the recommended dose of 2.5 mg/kg once monthly if there is no recurrence of severe or clinically significant transaminase elevations at the 1.25 mg/kg dose.

Renal Toxicity

Increases in serum creatinine levels and decreases in estimated glomerular filtration rate (eGFR) have been reported during treatment with GIVLAARI. In the placebo-controlled study, 15% of patients receiving GIVLAARI experienced a renally-related adverse reaction. The median increase in creatinine at Month 3 was 0.07 mg/dL. Monitor renal function during treatment with GIVLAARI as clinically indicated.

Injection Site Reactions

Injection site reactions were reported in 25% of patients receiving GIVLAARI in the placebo-controlled trial. Symptoms included erythema, pain, pruritus, rash, discoloration, or swelling around the injection site. One (2%) patient experienced a single, transient, recall reaction of erythema at a prior injection site with a subsequent dose administration.

Blood Homocysteine Increased

Increases in blood homocysteine levels have occurred in patients receiving GIVLAARI. In the ENVISION study, during the open label extension, adverse reactions of blood homocysteine increased were reported in 15 of 93 (16%) patients treated with GIVLAARI. Measure blood homocysteine levels prior to initiating treatment and monitor for changes during treatment with GIVLAARI. In patients with elevated blood homocysteine levels, assess folate, vitamins B12 and B6. Consider treatment with a supplement containing vitamin B6 (as monotherapy or a multivitamin preparation).

Drug Interactions

Concomitant use of GIVLAARI increases the concentration of CYP1A2 or CYP2D6 substrates, which may increase adverse reactions of these substrates. Avoid concomitant use of GIVLAARI with CYP1A2 or CYP2D6 substrates for which minimal concentration changes may lead to serious or life-threatening toxicities. If concomitant use is unavoidable, decrease the CYP1A2 or CYP2D6 substrate dosage in accordance with approved product labeling.

Adverse Reactions

The most common adverse reactions that occurred in patients receiving GIVLAARI were nausea (27%) and injection site reactions (25%).

**For additional information about GIVLAARI,
please see full Prescribing Information.**





Services Overview

Alnylam Assist™ offers a wide range of services to guide your patients through treatment with GIVLAARI® (givosiran)



How Alnylam Assist™ can help

After discussing treatment with your patient, begin the enrollment process by completing the Alnylam Assist™ Start Form. Upon receipt of the Start Form, an **Alnylam Case Manager dedicated to your patient's needs** will reach out to you and your patient within 2 business days.

Alnylam Assist™ will help with:

-  **Benefit verification**
-  **Education on the prior authorizations, claims, and appeals processes**
-  **Financial assistance program for eligible patients^a**
-  **Disease and product education**
-  **Ordering product for your patient**

Alnylam Field Reimbursement Directors (FRDs) are also available to you to provide education about the reimbursement process for Alnylam products.

FRDs will share their knowledge of:

- Billing and coding requirements for Alnylam products
- Chart documentation requirements
- Payer requirements


^a Patients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue the program at any time.

Please see Important Safety Information on page 3
and full Prescribing Information.





Personalized support for patients throughout the treatment process



Visit www.AlnylamAssist.com to complete the Start Form with your patient now.

^aPatients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue the program at any time.

Please see Important Safety Information on page 3 and full Prescribing Information.

 **GIVLAARI®**
(givosiran) injection for subcutaneous use
189 mg/mL



Helping your patients access Alnylam products

Benefit verification

Coverage for Alnylam products will vary by product, by plan, and by patient. Alnylam Assist™ can help determine patient-specific coverage requirements.

- To initiate a benefit verification for your patient, complete the **Start Form** electronically or via DocuSign at www.AlnylamAssist.com. You can also download and print the Start Form and fax it to 1-833-256-2747

Questions about how Alnylam Assist™ can help?

Call 1-833-256-2748

- Within 2 business days, your patient's dedicated Alnylam Case Manager will provide you and your patient with a benefit verification summary
- Alnylam Assist™ can provide information about patient financial assistance programs for eligible patients,^a if necessary (for additional information on financial assistance programs, see page 10)

^aPatients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue the program at any time.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





Treatment coverage

Alnylam Case Managers can explain the requirements and processes for prior authorizations, claims, and appeals.

Alnylam Assist™ can:

- Research the payer requirements as part of the benefit verification process
- Discuss the standard process for submitting a prior authorization and reimbursement claims
- Investigate reasons for denied or rejected prior authorizations, claims, and/or appeals

Alnylam's dedicated **Field Reimbursement Directors** are available to meet with you and your staff to answer coverage and reimbursement-related questions about Alnylam products

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.





Support for your patients

Alnylam Assist™ is here to help

Alnylam Assist™ will provide:

- An explanation of benefits so your patients understand their coverage
- Information about financial assistance programs for eligible patients^a
- A Patient Starter Kit, including educational materials designed to help patients understand their therapy and Alnylam Assist™
- Education for your patients from a designated Alnylam Patient Education Liaison (PEL)
 - Regionally based PELs are available to help patients gain a better understanding of the disease and treatment with Alnylam products
- Support throughout treatment with Alnylam product that is customized to each patient's communication preferences

^aPatients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue the program at any time.

**Please see Important Safety Information on page 3
and full Prescribing Information.**





Financial assistance programs^a

Patients may qualify for the following financial assistance programs^b:

- **Patient Assistance Program (PAP):** Provides GIVLAARI® (givosiran) at no cost to eligible patients, primarily the uninsured, who meet specified financial criteria.
- **Commercial Copay Program^c:** Covers certain out-of-pocket costs for eligible patients with commercial insurance

Eligibility criteria

PAP	Commercial Copay
Uninsured/functionally uninsured ^d	Commercially insured patients ^e
On-label diagnosis for prescribed Alnylam product	
	US residency (<i>including US territories</i>)
Financial eligibility requirements— supporting income documentation required ^e	Insurance must cover the prescribed Alnylam product

Once enrolled in Alnylam Assist™, an Alnylam Case Manager will review financial assistance programs your patient may qualify for based on his or her eligibility.

^aSome state laws may restrict or impact some aspects of these programs. An Alnylam Case Manager can provide additional information.

^bPatients must meet specified eligibility criteria to qualify for assistance. Alnylam reserves the right to make eligibility determinations and to modify or discontinue any program at any time.

^cPatients with Medicare, Medicaid, or other government-sponsored insurance are not eligible for the Alnylam Assist™ Commercial Copay Program. Out-of-pocket costs for the administration of GIVLAARI will not be covered for patients residing where it is prohibited by law or where otherwise restricted.

^dFunctionally uninsured patients are those who may be enrolled in a health plan but do not have coverage for an Alnylam product or cannot afford their cost share associated with their Alnylam product.

^eAcceptable forms of documentation may include: copy of most recent U.S. Individual Income Tax Return (IRS Form 1040); copy of most recent Social Security Benefit Statement (SSA-1099); copy of most recent pay stub. Patients with an income of ≤150% FPL are required to apply for Low Income Subsidy (LIS) for drugs filled through Medicare Part D.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





How to Complete the GIVLAARI® (givosiran) Start Form

Please see Important Safety Information on page 3 and full Prescribing Information.



How to complete the GIVLAARI® (givosiran) Start Form

This section will show you how to complete the Start Form. The notes on each page provide details to help ensure the form is filled out correctly. The Start Form serves as your patient's enrollment in Alnylam Assist™ and requires the signatures of both you and your patient. The Start Form also initiates your patient's prescription for GIVLAARI.

It is important to note the following before submitting the Start Form:

- Ensure highlighted key areas are correctly filled out
- Confirm that you and your patient sign where indicated

Options for getting started

1. Complete and submit the **electronic Start Form** with your patient **or**
2. Complete the **paper Start Form** with your patient and fax to 1-833-256-2747 **or**
3. Begin the Start Form, filling in all details needed by a healthcare professional, and then have your patient complete the form via **DocuSign**



All 3 options to get started can be found at www.AlnylamAssist.com.

**Please see Important Safety Information on page 3
and full Prescribing Information.**





For patients

Your Patient's Email

Please make sure your patients fill in this field.

Preferred Phone Number & Voicemail Checkbox

By allowing Alnylam Assist™ to leave voicemails, delays in benefit verification and other communications can be avoided.

Language Translation

Alnylam Assist™ offers translation services for non-English-speaking patients.

Insurance Information

Patients (or their authorized representatives) can fill in the provided fields or attach copies of both sides of their insurance and pharmacy benefits cards.

Start Form

For Patients
Alnylam Assist® Enrollment
(Sections 1 – 5 to be read and completed by **Patient or Patient's Authorized Representative**)

The purpose of this form is to permit Alnylam Assist® participants to receive information and support ("Patient Support") from Alnylam Pharmaceuticals, Inc., its affiliates, representatives, agents, and contractors ("Alnylam"). Alnylam Assist® provides Patient Support to eligible patients who have been prescribed an Alnylam medicine. This includes: (1) providing reimbursement and financial support to eligible patients (such as investigating your insurance coverage, confirming out-of-pocket costs, and reviewing eligibility for financial assistance); (2) working with you and your provider to fill your prescription; and (3) providing you with disease and medication-related educational resources and communications.

Please read this form carefully and ask any questions that you may have before signing.

1. Patient Information

Name (First, Mi, Last): Lawrence N. Reele	Date of Birth: Month/Day/Year 05/14/1956	Email: LNReele@email.com
Street Address: 1020 Generic Ave.		City: Springfield
		State: MA
		ZIP: 15123
Preferred Phone Number: <input checked="" type="checkbox"/> Okay to leave voicemail (555) 137-1634		Alternative Phone Number (if available): <input type="checkbox"/> Okay to leave voicemail (555) 136-1522
Caregiver Name (optional): Diane Reele		Caregiver Relationship to Patient (optional): Wife
Caregiver Phone (optional): (555) 136-1522		
Language translation? <input checked="" type="checkbox"/> Yes, translation needed <input type="checkbox"/> No If yes, please indicate language: Portuguese		

2. Insurance Information Attach a copy of both sides of your INSURANCE and PRESCRIPTION cards Check if you do not have insurance

Primary Insurance Provider: ABC Insurance Co.	Employer Name: Company Inc.	Policy Number: 123456789101	Group Number: 12-34567
Policyholder Name (First, Mi, Last), if other than the patient: 		Policyholder Date of Birth: Month/Day/Year 	
Insurance Phone: (555) 136-2222			
Pharmacy Plan Provider (if applicable): 	Policy Number: 	Group Number: 	Rx BIN Number:
Policyholder Name (First, Mi, Last), if other than the patient: 		Policyholder Date of Birth: Month/Day/Year 	
Secondary Insurance Provider (if applicable): 	Employer Name: 	Policy Number: 	Group Number:
Policyholder Name (First, Mi, Last), if other than the patient: 		Policyholder Date of Birth: Month/Day/Year 	
Insurance Phone: 			

> Continue to page 2 to complete the patient portion of the Start Form

Please see Important Safety Information on page 4, and full Prescribing Information available from your Alnylam representative or at www.givlaari.com.

— 1 of 4 —

Please see Important Safety Information on page 3 and full Prescribing Information.

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL

13



Authorization to share protected health information/authorization for Alnylam Assist™ enrollment

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL

Start Form

Alnylam ASSIST

3. Authorization to Share Protected Health Information

By signing below, I authorize my healthcare providers, including my physicians and pharmacies ("My Providers") and my health insurance plan ("My Plan") to share my medical information (such as information about my diagnosis, prescriptions, and treatment) and my insurance information ("My Information") with Alnylam so that Alnylam can provide Patient Support. I authorize My Providers to use My Information to provide me with certain offerings related to my medication and treatment. I understand that my pharmacy will receive payment from Alnylam for disclosing My Information to Alnylam. I understand that once My Information has been disclosed, federal privacy laws may no longer protect the information. However, I understand that Alnylam agrees to protect My Information by using and disclosing it only for purposes described in this Authorization or as required by law or regulations. I understand that I may refuse to sign this Authorization, and that my treatment, insurance enrollment, and eligibility for insurance benefits are not conditioned upon signing this Authorization. I also understand, however, that refusing to sign this Authorization means that I may not participate in Alnylam Assist® and may not be able to take advantage of other offerings by Alnylam. I may cancel or revoke this Authorization at any time by mailing a letter to Privacy Officer at Alnylam, Attn: Legal Department, 675 West Kendall Street, Cambridge, MA 02142 or by sending an email to privacy@alnylam.com. I understand that if I revoke this Authorization, My Providers and Alnylam will stop using and sharing My Information under this Authorization, but my revocation will not affect uses and disclosures of My Information prior to my revocation in reliance upon this Authorization.

This Authorization expires ten (10) years from the date signed below, or earlier if required by state or local law, unless I revoke it before then. I understand that I may receive a copy of this Authorization.

Lawrence N. Reece	X 
Print Patient or Authorized Patient Representative Name	Signature of Patient or Authorized Patient Representative
Relationship to Patient	January 1, 2021
Date	

4. Authorization for Alnylam Assist® and Communications

By signing below, I confirm I would like to enroll in the Alnylam Assist® program and authorize Alnylam to provide me with Patient Support. I understand that Alnylam Assist® is an optional program.

I agree that Alnylam may use My Information and share it with My Providers or My Plan in connection with providing the Patient Support, administering the Alnylam Assist® program, or as otherwise required by Alnylam to meet its legal obligations. For example, Alnylam may communicate with me (such as by mail, phone, email, and/or text message) or my caregiver, use My Information to tailor the Alnylam Assist®-related communications to my needs, and share information with My Providers about dispensing my Alnylam medicine to me. I understand that Alnylam may de-identify My Information, combine it with information about other patients, and use the resulting information for Alnylam's business purposes.

Lawrence N. Reece	X 
Print Patient or Authorized Patient Representative Name	Signature of Patient or Authorized Patient Representative
Relationship to Patient	January 1, 2021
Date	

5. Opt In to Receive Marketing Communications (optional)

By checking this box, I authorize Alnylam, and companies working with Alnylam, to contact me by mail, email, fax, and/or telephone regarding marketing and promotional communications, customer surveys, or for market research surveys. **I understand that I am not required to provide this consent as a condition of receiving any Alnylam medicine or services from Alnylam.**

Please see **Important Safety Information** on page 4, and full **Prescribing Information** available from your Alnylam representative or at www.givlaari.com.

— 2 of 4 —

**Please see Important Safety Information on page 3
and full Prescribing Information.**

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL



For healthcare providers

Product Acquisition

Select your preferred method of product acquisition (specialty pharmacy or specialty distributor). If acquisition method is unknown, select *Unknown*.

GIVLAARI® (givosiran) Dosing Information

- Confirm that your patient is being prescribed GIVLAARI as indicated by **checking the box**
- Make sure to include the **primary diagnosis code** and **patient's weight (kg)**

GIVLAARI Prescription

Ensure you fill in this prescription field for your patients.

Signature of Prescriber

Prescriber should only sign one prescription field and include date in Section 7.

To prevent a generic substitution, sign the "dispense as written" field.

To allow generic substitutions, sign the "substitution permitted" field.



Start Form



For Healthcare Providers

(Sections 6 – 8 to be read and completed by **Healthcare Provider**)

6. Prescriber Information

Name (First, Last): Charles Sample	Sample Co.	Practice Name: Neurology	Specialty:
Practice Street Address: 530 Pioneer Road	City: Easton	State: MA	
ZIP: 40520	Phone: (555) 876-5309	Fax:	National Provider ID (NPI) #: 1234567890
Office Contact Name: Jane Smith	Phone: (555) 652-5678		State License #: S943072
Email: SampleDoc@email.com			Anticipated First Treatment Date: January 1, 2021
Product acquisition: <input checked="" type="checkbox"/> Specialty Pharmacy <input checked="" type="checkbox"/> Accredo <input type="checkbox"/> PANTHERx <input type="checkbox"/> No preference <input type="checkbox"/> Specialty Distributor (McKesson Specialty or McKesson Plasma and Biologics) <input type="checkbox"/> Unknown			

7. GIVLAARI®(givosiran) Prescription

(This is a prescription; a prescriber's signature and date are required.)

Patient Name:	Patient Date of Birth: Month/Day/Year:				
<input checked="" type="checkbox"/> E80.20 (Unspecified porphyria) <input checked="" type="checkbox"/> E80.21 (Acute intermittent (hepatic) porphyria) <input type="checkbox"/> E80.29 (Other porphyria) <input type="checkbox"/> Other					
GIVLAARI Injection for subcutaneous use, 189 mg/mL <small>(Recommended dose is 2.5 mg/kg monthly)</small>					
Date Patient Weight Taken	Patient Weight (in kg)	Total Calculated Dose (SC monthly)	Number of Vials/Treatment	Refills	
1/1/2021	61.23	(mg) 153.07 (mL)	1	189 mg/mL vial(s)	<input checked="" type="checkbox"/> Refill x 11 <input type="checkbox"/> Other
Any known allergies? <input type="checkbox"/> Yes <input checked="" type="checkbox"/> No If yes, please list: Oxycodeone					
List or attach a list of concomitant medications: Oxycodeone					
Special Instructions: None					
<input type="checkbox"/> If acquiring through Accredo or PANTHERx, please check here to authorize ancillary supplies, such as needles and syringes, as needed to administer treatment. <input type="checkbox"/> I confirm that my patient is being prescribed GIVLAARI for the treatment of acute hepatic porphyria (AHP) in adults.					

I authorize Alnylam to act on my behalf for the limited purposes of transmitting this prescription to the appropriate pharmacy.
I will comply with my state-specific prescription requirements, such as e-prescribing, state-specific prescription form, fax language, etc.

X Charles	Date January 1, 2021
Prescriber Signature (No Stamps) Dispense as Written	Date
X	Date
Prescriber Signature (No Stamps) Substitution Permitted	
Desired Site of Care <input type="checkbox"/> Home Injection (see patient home address) <input type="checkbox"/> Alternate Medical Facility (provide facility name and address) Facility Name/Address _____	
<input checked="" type="checkbox"/> Physician Office (see provider office address) <input type="checkbox"/> Facility to Home (first dose at facility; remainder at home)	

► Continue to page 4 to complete the prescriber portion of the Start Form

Please see **Important Safety Information** on page 4, and full **Prescribing Information** available from your Alnylam representative or at www.givlaari.com.

— 3 of 4 —

Desired Site of Care

Ask your patient where he or she would like to receive treatment.

Please see Important Safety Information on page 3 and full Prescribing Information.





Prescriber declaration

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL

Alnylam ASSIST™

8. Prescriber Declaration

By signing below, I certify that:

- ▷ The information contained in this form is complete and accurate to the best of my knowledge.
- ▷ I understand that Alnylam is not responsible for filing claims or submitting other information to my patient's insurer and that the information provided by Alnylam Assist® is educational in nature.
- ▷ I understand that my patient may authorize Alnylam Assist® to provide Patient Support. I understand that this program does not include individual treatment or medical advice to the patient, and it does not replace the medical treatment and care provided by me as the patient's healthcare provider.
- ▷ I further certify that I understand that any support provided by Alnylam Assist® on behalf of any patient is not made in exchange for any express or implied agreement or understanding that I would recommend, prescribe, or use GIVLAARI® (givosiran) or any other Alnylam product, and any decision to prescribe GIVLAARI was, and in the future will be, based solely on my determination of medical necessity.
- ▷ I have obtained the required authorizations from my patient to release the referenced medical and/or other patient information relating to my patient's treatment to Alnylam Assist®.

 Prescriber signature (stamps not acceptable)

January 1, 2021 Date

INDICATION
GIVLAARI® (givosiran) is indicated for the treatment of adults with acute hepatic porphyria (AHP).

IMPORTANT SAFETY INFORMATION

Contraindications
GIVLAARI is contraindicated in patients with known severe hypersensitivity to givosiran. Reactions have included anaphylaxis.

Anaphylactic Reaction
Anaphylaxis has occurred with GIVLAARI treatment (<1% of patients in clinical trials). Ensure that medical support is available to appropriately manage anaphylactic reactions when administering GIVLAARI. Monitor for signs and symptoms of anaphylaxis. If anaphylaxis occurs, immediately discontinue administration of GIVLAARI and institute appropriate medical treatment.

Hepatic Toxicity
Transaminase elevations (ALT) of at least 3 times the upper limit of normal (ULN) were observed in 15% of patients receiving GIVLAARI in the placebo-controlled trial. Transaminase elevations primarily occurred between 3 to 5 months following initiation of treatment. Measure liver function tests prior to initiating treatment with GIVLAARI, repeat every month during the first 6 months of treatment, and as clinically indicated thereafter. Interrupt or discontinue treatment with GIVLAARI for severe or clinically significant transaminase elevations. In patients who have dose interruption and subsequent improvement, reduce the dose to 1.25 mg/kg once monthly. The dose may be increased to the recommended dose of 2.5 mg/kg once monthly if there is no recurrence of severe or clinically significant transaminase elevations at the 1.25 mg/kg dose.

Renal Toxicity
Increases in serum creatinine levels and decreases in estimated glomerular filtration rate (eGFR) have been reported during treatment with GIVLAARI. In the placebo-controlled study, 15% of patients receiving

GIVLAARI experienced a renally-related adverse reaction. The median increase in creatinine at Month 3 was 0.07 mg/dL. Monitor renal function during treatment with GIVLAARI as clinically indicated.

Injection Site Reactions
Injection site reactions were reported in 25% of patients receiving GIVLAARI in the placebo-controlled trial. Symptoms included erythema, pain, pruritus, rash, discoloration, or swelling around the injection site. One (2%) patient experienced a single, transient, recall reaction of erythema at a prior injection site with a subsequent dose administration.

Blood Homocysteine Increased
Increases in blood homocysteine levels have occurred in patients receiving GIVLAARI. In the ENVISION study, during the open label extension, adverse reactions of blood homocysteine increased were reported in 15 of 93 (16%) patients treated with GIVLAARI. Measure blood homocysteine levels prior to initiating treatment and monitor for changes during treatment with GIVLAARI. In patients with elevated blood homocysteine levels, assess folate, vitamins B12 and B6. Consider treatment with a supplement containing vitamin B6 (as monotherapy or a multivitamin preparation).

Drug Interactions
Concomitant use of GIVLAARI increases the concentration of CYP1A2 or CYP2D6 substrates, which may increase adverse reactions of these substrates. Avoid concomitant use of GIVLAARI with CYP1A2 or CYP2D6 substrates for which minimal concentration changes may lead to serious or life-threatening toxicities. If concomitant use is unavoidable, decrease the CYP1A2 or CYP2D6 substrate dosage in accordance with approved product labeling.

Adverse Reactions
The most common adverse reactions that occurred in patients receiving GIVLAARI were nausea (27%) and injection site reactions (25%).

For additional information about GIVLAARI, please see full Prescribing Information available from your Alnylam representative or at givlaari.com.

Fax the completed Start Form to 1-833-256-2747 | Call Alnylam Assist® at 1-833-256-2748 8AM–6PM ET, Monday–Friday | For more information, visit www.AlnylamAssist.com

Alnylam® PHARMACEUTICALS GIVLAARI and Alnylam Assist are registered trademarks of Alnylam Pharmaceuticals, Inc. © 2021 Alnylam Pharmaceuticals, Inc. All rights reserved. AS1-USA-00102-V3

— 4 of 4 —

Signature of Prescriber


Sign and date the declaration on the last page, certifying the information provided in the form and authorization of services. Before submitting the form, ensure **both prescriber signatures** are provided in Sections 7 and 8.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL



Once the completed Start Form is received by Alnylam Assist™



^aIf a reauthorization is required, a new request must be submitted.

^bAlnylam Assist™ can provide education on prior authorization requirements and processes, but cannot guarantee that a patient's prior authorization will be approved.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.





Overview of Acquisition Process for GIVLAARI® (givosiran)

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.



Overview of acquisition process

1. Identify a medically appropriate patient for GIVLAARI® (givosiran)

- GIVLAARI is indicated for the treatment of adults with acute hepatic porphyria (AHP)

2. Verify insurance benefits

- Submit Start Form to Alnylam Assist™
- Within 2 business days after Start Form submission, an Alnylam Case Manager will reach out to you and your patient. Following a benefit investigation, you will receive a copy of your patient's Summary of Benefits and Coverage, including prior authorization and payer requirements, as well as financial program information for eligible patients
- Coverage for GIVLAARI will vary by plan and by patient

3. Obtain GIVLAARI

- Specialty Distributor: **McKesson**—healthcare professional can order from either McKesson Specialty Health or McKesson Plasma and Biologics
- Specialty Pharmacy: either **Accredo** or **PANTHERx** will coordinate drug shipment with healthcare professional

NOTE: GIVLAARI is obtained via a limited network of distributors highlighted above.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





Billing and Coding Overview

Please see Important Safety Information on page 3 and full Prescribing Information.



Coverage, coding, and payment

GIVLAARI® (givosiran) received U.S. Food and Drug Administration (FDA) approval on November 20, 2019, and is indicated for the treatment of adults with acute hepatic porphyria (AHP).

Coverage

- **For Medicare patients** receiving GIVLAARI who are covered under Medicare Part B, the Medicare Administrative Contractors (MACs) may require additional documentation to determine the medical necessity of the treatment, although prior authorization is not required^{a,b}
- **For patients enrolled in a State Medicaid or commercial health plan**, GIVLAARI coverage will vary by payer^a

Payment

Payer Type	Payment Methodology
Medicare Fee-for-Service	Average Sales Price (ASP) + 6% ^c
State Medicaid and Commercial Payers	Payment rates will vary by payer and provider contract

^aIt is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for products and services rendered. Providers should contact third-party payers for specific information on their coding, coverage, and payment policies.

^bMedicare Advantage Plans may require a prior authorization for GIVLAARI.

^cDoes not account for any required payment reductions if sequestration is in effect.

Alnylam **Field Reimbursement Directors** are available to meet with you and your staff to answer coverage, coding and payment questions about GIVLAARI. Contact Alnylam Assist™ at 1-833-256-2748.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.





Billing and Coding

Physician Office

Please see Important Safety Information on page 3 and full Prescribing Information.



Coding

Please refer to the table below to support appropriate claims submission for GIVLAARI® (givosiran).

Code Type	Code	Code Description
ICD-10-CM	E80.20	Unspecified porphyria
	E80.21	Acute intermittent (hepatic) porphyria
	E80.29	Other porphyria
CPT® ^a	96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular
HCPCS	J0223	Injection, givosiran, 0.5 mg
NDC	10-digit: 71336-1001-1 11-digit: 71336-1001-01	189 mg/mL single-dose vial

^aCPT Copyright 2019 American Medical Association. All rights reserved.

CPT® is a registered trademark of the American Medical Association.

Applicable FARS/DFARS Restrictions Apply to Government Use.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

ICD-10-CM=International Classification of Diseases, 10th Revision, Clinical Modification; CPT=Current Procedural Terminology; HCPCS=Healthcare Common Procedure Coding System; NDC=National Drug Code.

Providers should consult the ICD-10-CM code book and use their own clinical judgment to confirm coding.

Please see Important Safety Information on page 3 and full Prescribing Information.





Physician office: sample CMS-1500 claim form

GIVLAARI® (givosiran) and the associated services provided in a physician office are billed on the CMS-1500 claim form or its electronic equivalent. A sample CMS-1500 claim form for billing GIVLAARI is provided on the next page.

- The following CMS-1500 claim form for GIVLAARI is for illustrative purposes
- It is the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for the products and services rendered
- Providers should contact payers for specific information on their coding, coverage, and payment policies
- Some payers may require the use of a JW modifier when billing for the unused portion of the single-dose vial (wastage)—providers should contact payers about specific coding and payment policies
- Providers should contact their billing software vendors to ensure that they are utilizing the recommended loops and segments

Dosing calculation example

- GIVLAARI is supplied as a 189 mg/mL solution in a single-dose vial
- The recommended dose of GIVLAARI is 2.5 mg/kg administered via subcutaneous injection once monthly
- Dosing is based on actual body weight

Calculation

How to Calculate Dosage (mg)	How to Calculate Injection Volume (mL)
(body weight [kg] × 2.5 mg/kg) = mg	(mg × 1 mL/189 mg) = mL

Example - 68 kg patient

Dosage (mg)	Injection Volume (mL)
68 kg × 2.5 mg/kg = 170 mg	170 mg × 1 mL/189 mg = 0.9 mL

Please see Important Safety Information on page 3
and full Prescribing Information.





LOCATOR

21

Enter the appropriate primary diagnosis code from the patient's medical record in Locator 21A.

LOCATOR

21 ICD-IND

Enter "0" to indicate use of ICD-10-CM diagnosis coding system.

LOCATOR

24 A-B

Enter the date of service and the appropriate place of service code.

LOCATOR

24

Enter the GIVLAARI® (givosiran) HCPCS code J0223 on the first line and the CPT code 96372 for drug administration on the second line.

Sample CMS-1500 Claim Form

The form includes sections for Carrier information, Patient and Insured Information, Physician or Supplier Information, and a large section for Item Details (Lines 1-6). Arrows point from the locator numbers to specific fields on the form:

- Locator 21:** Points to the Primary Diagnosis field (Item 1).
- Locator 21 ICD-IND:** Points to the ICD-10-CM indicator field (Item 1).
- Locator 24 A-B:** Points to the Date of Service (Item 14) and Place of Service (Item 15) fields.
- Locator 24:** Points to the HCPCS code J0223 in the first line item and the CPT code 96372 in the second line item.

LOCATOR

24D

Enter the appropriate HCPCS code for GIVLAARI: J0223 (injection, givosiran, 0.5 mg).

Shaded area of Locator 24D (when applicable): N4713361000101 MLX (X = number of vials; for example, ML1 = 1 vial, ML2 = 2 vials, etc.)

LOCATOR

24E

Specify the diagnosis, from Locator 21, that relates to the product or procedure listed in Locator 24d.

LOCATOR

24G

Enter the number of service units for each line item.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL



Clean claim filing checklist



Select the appropriate primary diagnosis



Confirm appropriate clinical documentation to support diagnosis



Understand any payer-specific requirements (prior authorization, coding details, etc)



Utilize all appropriate ICD-10, CPT®, and HCPCS codes

- For all claims in the physician office setting, use HCPCS J0223 (Injection, givosiran, 0.5 mg) for dates of services on or after July 1, 2020^a
 - Remember: Billing Unit = 0.5 mg
- Remember to use the sample claim form on page 25 as a guide



Anticipate requests from payers for additional clinical information prior to claims being processed for payment

It is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for those products and services rendered. Contact third-party payers for specific information on their coding and payment policies.

^aHCPCS codes for GIVLAARI® (givosiran) may vary for dates of service prior to July 1, 2020.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





Billing and Coding

Hospital Outpatient Department

Please see Important Safety Information on page 3 and full Prescribing Information.



Coding

Please refer to the table below to support appropriate claims submission for GIVLAARI® (givosiran).

Code Type	Code	Code Description
ICD-10-CM	E80.20	Unspecified porphyria
	E80.21	Acute intermittent (hepatic) porphyria
	E80.29	Other porphyria
CPT® ^a	96372	Therapeutic, prophylactic, or diagnostic injection (specify substance or drug); subcutaneous or intramuscular
HCPCS	J0223	Injection, givosiran, 0.5 mg
	0250	General pharmacy
Revenue	0940	Other therapeutic services
	0636	Drugs requiring detailed coding
NDC	10-digit: 71336-1001-1 11-digit: 71336-1001-01	189 mg/mL single-dose vial

^aCPT Copyright 2019 American Medical Association. All rights reserved.

CPT® is a registered trademark of the American Medical Association.

Applicable FARS/DFARS Restrictions Apply to Government Use.

Fee schedules, relative value units, conversion factors and/or related components are not assigned by the AMA, are not part of CPT, and the AMA is not recommending their use. The AMA does not directly or indirectly practice medicine or dispense medical services. The AMA assumes no liability for data contained or not contained herein.

ICD-10-CM=International Classification of Diseases, 10th Revision, Clinical Modification; CPT=Current Procedural Terminology; HCPCS=Healthcare Common Procedure Coding System; NDC=National Drug Code.

Providers should consult the ICD-10-CM code book and use their own clinical judgment to confirm coding.

Please see Important Safety Information on page 3 and full Prescribing Information.





Hospital outpatient: sample UB-04 claim form

GIVLAARI® (givosiran) and the associated services provided in a hospital outpatient department setting are billed on the UB-04 claim form or its electronic equivalent. A sample UB-04 claim form for billing GIVLAARI is provided on the next page.

- The following UB-04 claim form for GIVLAARI is for illustrative purposes
- It is the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for the products and services rendered
- Providers should contact payers for specific information on their coding, coverage, and payment policies
- Some payers may require the use of a JW modifier when billing for the unused portion of the single-dose vial (wastage)—providers should contact payers about specific coding and payment policies
- Providers should contact their billing software vendors to ensure they are utilizing the recommended loops and segments

Dosing calculation example

- GIVLAARI is supplied as a 189 mg/mL solution in a single-dose vial
- The recommended dose of GIVLAARI is 2.5 mg/kg administered via subcutaneous injection once monthly
- Dosing is based on actual body weight

Calculation

How to calculate Dosage (mg)	How to calculate Injection Volume (mL)
(body weight [kg] × 2.5 mg/kg) = mg	(mg × 1 mL/189 mg) = mL

Example - 68 kg patient

Dosage (mg)	Injection Volume (mL)
68 kg × 2.5 mg/kg = 170 mg	170 mg × 1 mL/189 mg = 0.9 mL

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





LOCATOR

42

List the appropriate revenue code for the service provided.

For Medicare, use the revenue code
0636—Drugs requiring detailed coding.

For payers other than Medicare, the revenue code for GIVLAARI may vary, although some private payers and **Medicaid** plans accept revenue code 0250—General pharmacy.

LOCATOR

43

Enter the corresponding description for the revenue code listed in Locator 42 **OR** Enter the NDC (when applicable): N471336100301 MLX (X = number of vials; for example, ML1 = 1 vial, ML2 = 2 vials, etc.)

LOCATOR

44

Enter the appropriate HCPCS code for GIVLAARI: J0223 (injection, givosiran, 0.5 mg).

Sample UB-04 Claim Form

1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	
PATIENT NAME										PATIENT ADDRESS										BILLING INFORMATION																													
b										c										d																													
10 BIRTHDATE		11 SEX		12 DATE		ADMISSION		13 HR		14 TYPE		15 SRC		16 DHR		17 STAT		18		19		20		21		CONDITION CODES		22		23		24		25		26		27		28		29 AGO/STATE		30					
31 OCCURRENCE DATE		32 OCCURRENCE DATE		33 OCCURRENCE DATE		34 OCCURRENCE DATE		35 OCCURRENCE DATE		36 OCCURRENCE SPAN FROM		37 OCCURRENCE SPAN THROUGH		38		39		40		41		42		43		44		45		46		47		48		49													
CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE									
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d											
42 REV CD		43 DESCRIPTION		44 HOPCS / RATE / HCPCS CODE		45 SERV. DATE		46 SERV. UNITS		47 TOTAL CHARGES		48 NON-COVERED CHARGES		49		50		51		52		53		54		55		56		57		58		59		60		61		62									
PAGE		OF		CREATION DATE		TOTALS																																											
50 PAYER NAME		51 HEALTH PLAN ID		52 REL INFO		53 AGO/STATE		54 PRIOR PAYMENTS		55 EST. AMOUNT DUE		56 NPI		57		58		59		60		61		62		63		64		65		66		67		68		69		70		71		72		73			
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d			
58 INSURED'S NAME		59 PEL		60 INSURED'S UNIQUE ID		61 GROUP NAME		62 INSURANCE GROUP NO.		63 TREATMENT AUTHORIZATION CODES		64 DOCUMENT CONTROL NUMBER		65 EMPLOYER NAME		66		67		68		69		70		71		72		73		74		75		76		77		78		79		80					
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d			
65		66		67		68		69		70		71		72		73		74		75		76		77		78		79		80		81		82		83		84		85		86		87		88		89	
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d			
69 ADMT DX		70 PATIENT REASON DX		71 OTHER PROCEDURE		72 OTHER PROCEDURE		73		74		75		76		77		78		79		80		81		82		83		84		85		86		87		88		89									
CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE		CODE							
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d											
80 REMARKS		81		82		83		84		85		86		87		88		89		90		91		92		93		94		95		96		97		98		99											
a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d		a		b		c		d											

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.



LOCATOR

40

Enter the GIVLAARI®(givosiran) HCPCS code J0223 on the first line and the CPT code 96372 for drug administration on the second line.

LOCATOR

45

Enter the service date.

LOCATOR

46

Enter the number of service units for each line item.

LOCATOR

47

Enter the total charge for each line item.

LOCATOR

66

Enter “0” to indicate use of the ICD-10-CM diagnosis coding system.

LOCATOR

67

Enter the primary diagnosis code.



Clean claim filing checklist



Select the appropriate primary diagnosis



Confirm appropriate clinical documentation to support diagnosis



Understand any payer-specific requirements (prior authorization, coding details, etc)



Utilize all appropriate ICD-10, CPT®, HCPCS, and Revenue codes

- For all claims in the hospital outpatient department setting, use HCPCS code J0223 (Injection, givosiran, 0.5 mg) for dates of service on or after July 1, 2020^a
 - Remember: Billing Unit = 0.5 mg
- Remember to use the sample claim form on page 30 as a guide



Anticipate requests from payers for additional clinical information prior to claims being processed for payment

It is always the provider's responsibility to determine the appropriate healthcare setting and to submit true and correct claims for those products and services rendered. Contact third-party payers for specific information on their coding and payment policies

^aHCPCS codes for GIVLAARI® (givosiran) will vary for dates of service prior to July 1, 2020.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





Copay Claim Submission Guide

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.



Copay claim submission



The following outline will show you how to submit a medical benefits claim, pharmacy claim, or mail-in rebate for GIVLAARI® (givosiran). Before submitting a claim, please ensure the following:

- The patient is enrolled in Alnylam Assist™ (via the Start Form found at AlnylamAssist.com)
- The patient's benefits have been verified
- The patient has provided his or her medical benefit or pharmacy member number


Upon a patient's enrollment into the copay program, an Alnylam Case Manager will provide your practice with the patient's Payer ID, Group Number, and Member Number required to submit a copay claim.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.





Submitting copay claims for GIVLAARI® (givosiran)



^aMedical Benefits or Pharmacy Claims.

TrialCard® is a registered trademark of TrialCard Incorporated.

**Please see Important Safety Information on page 3
and full Prescribing Information.**

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL



Dosing and Administration Guide

INDICATION

GIVLAARI® (givosiran) is indicated for the treatment of adults with acute hepatic porphyria (AHP).

IMPORTANT SAFETY INFORMATION

Contraindications

GIVLAARI is contraindicated in patients with known severe hypersensitivity to givosiran. Reactions have included anaphylaxis.



Overview of GIVLAARI® (givosiran)

GIVLAARI is indicated for the treatment of adults with acute hepatic porphyria (AHP).

GIVLAARI is a double-stranded RNAi therapeutic specifically targeting *ALAS1* mRNA, reducing *ALAS1* mRNA levels and leading to reductions in urinary ALA and PBG.

GIVLAARI dosing¹

Weight-based dosing with GIVLAARI

The recommended dose of GIVLAARI is 2.5 mg/kg administered via subcutaneous injection once monthly. Dosing is based on actual body weight.

Dose Modification for Adverse Reactions:

In patients with severe or clinically significant transaminase elevations, who have dose interruption and subsequent improvement, reduce the dose to 1.25 mg/kg once monthly. In patients who resume dosing at 1.25 mg/kg once monthly without recurrence of severe or clinically significant transaminase elevations, the dose may be increased to the recommended 2.5 mg/kg once monthly.

The following chart contains several illustrative dosing calculations. These examples are not intended to replace your clinical judgment. Confirm all calculations prior to GIVLAARI administration.

- GIVLAARI is a ready-to-use solution that does not require additional reconstitution or dilution prior to administration, supplied in single-dose vials of 189 mg/mL
- Calculate volume required based on recommended dosage: patient weight in kg × 2.5 mg/kg × 1 mL/189 mg = mL of GIVLAARI to administer
- If the total volume of GIVLAARI per dose is >1.5 mL, divide the dose into multiple injections of approximately equal volumes

DOSING BY WEIGHT				
GIVLAARI DOSE	BODY WEIGHT (kg)	TOTAL DOSE VOLUME* (mL)	NUMBER OF VIALS	INJECTION(S)
2.5 mg/kg	40	0.5		1 injection in a 1-mL syringe
	50	0.7		
	60	0.8		
	70	0.9		
	80	1.1		1 injection in a 3-mL syringe
	90	1.2		
	100	1.3		
	110	1.5		
120	1.6 [†]			2 injections: 1. 0.8 mL in a 1-mL syringe 2. 0.8 mL in a 1-mL syringe

*Volumes rounded to nearest 0.1 mL (eg, 0.55 mL was rounded to 0.6 mL).

[†]Divide doses requiring volumes >1.5 mL equally into multiple syringes.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.






GIVLAARI® (givosiran) administration instructions¹

- Ensure that medical support is available to appropriately manage anaphylactic reactions when administering GIVLAARI
- GIVLAARI is intended for subcutaneous use by a healthcare professional only

Supplies needed



Product vial



Sterile syringe(s)
(1 mL or 3 mL)



21G or larger
needle to
withdraw product
from vial



25G or 27G needle
with 1/2" or 5/8" length
for subcutaneous
administration




Alcohol swabs




Sharps
container for
needle disposal

Preparing the GIVLAARI dose²

- Use aseptic technique
- Calculate the required volume of GIVLAARI based on the recommended weight-based dosage
- GIVLAARI is a sterile, preservative-free, clear, colorless-to-yellow solution. As with all parenteral drug products, inspect the solution for particulate matter and discoloration prior to administration



1. To withdraw GIVLAARI solution, hold the vial upright or tilt at a slight angle and ensure the flat edge of the needle is pointed downwards
2. Gently draw up the necessary volume(s) of GIVLAARI solution with a 21G or larger needle into either:
 - A sterile 1-mL syringe for volumes up to 1 mL or
 - A sterile 3-mL syringe for volumes >1 mL, up to a maximum of 1.5 mL
3. Point the needle and syringe straight upwards and tap the syringe to move any air bubbles to the top
4. Once the air bubbles are at the top, gently push the plunger to expel the bubbles from the syringe
 - Check to make sure you still have the correct amount of GIVLAARI in the syringe
5. Once the dose is prepared and in the administration syringe(s), replace the 21G or larger needle with either a 25G or 27G needle
 - **Note: Do not advance GIVLAARI into the 25G or 27G needle**



IMPORTANT SAFETY INFORMATION

Anaphylactic Reaction


Anaphylaxis has occurred with GIVLAARI treatment (<1% of patients in clinical trials). Ensure that medical support is available to appropriately manage anaphylactic reactions when administering GIVLAARI. Monitor for signs and symptoms of anaphylaxis. If anaphylaxis occurs, immediately discontinue administration of GIVLAARI and institute appropriate medical treatment.

Please see Important Safety Information on page 3 and full Prescribing Information.





Choosing and preparing the injection site¹



- You may inject GIVLAARI® (givosiran) into the abdomen, the back or side of the upper arms, or the thighs
 - Rotate injection sites
- **Note:**
 - When administering subcutaneous injections into the abdomen, avoid a 5.0-cm diameter circle around the navel
 - If more than one injection is needed for a single dose of GIVLAARI, the injection sites should be at least 2 cm apart from previous injection locations
- Clean the area you intend to inject with an alcohol swab and wait for the area to dry completely

DO NOT ADMINISTER INJECTIONS¹:

- In areas that are **reddened, inflamed, or swollen**
- Into **scar tissue**

IMPORTANT SAFETY INFORMATION

Hepatic Toxicity

Transaminase elevations (ALT) of at least 3 times the upper limit of normal (ULN) were observed in 15% of patients treated with GIVLAARI in the placebo-controlled trial. Transaminase elevations primarily occurred between 3 to 5 months following initiation of treatment.



Measure liver function tests prior to initiating treatment with GIVLAARI, repeat every month during the first 6 months of treatment, and as clinically indicated thereafter. Interrupt or discontinue treatment with GIVLAARI for severe or clinically significant transaminase elevations. In patients who have dose interruption and subsequent improvement, reduce the dose to 1.25 mg/kg once monthly. The dose may be increased to the recommended dose of 2.5 mg/kg once monthly if there is no recurrence of severe or clinically significant transaminase elevations at the 1.25 mg/kg dose.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.

 **GIVLAARI®**
(givosiran) injection for subcutaneous use
189 mg/mL



Administering GIVLAARI® (givosiran)²



1. Ensure proper injection technique. Do not inject into a vein or muscle
2. With your free hand, pinch the skin at the injection site and insert the needle at a right angle (90 degrees) to deliver the injection just below the skin
 - In patients with little subcutaneous tissue or if the needle size is longer than 1 inch, the needle should be inserted at a 45-degree angle
 - Do not press down on the plunger while piercing the skin
3. Once the needle is inserted through the skin, release the pinched skin and administer the dose in a slow and steady manner
 - **Note: Do not aspirate after inserting the needle to prevent tissue damage, hematoma, and bruising**
4. Once GIVLAARI has been administered, count for at least 5 seconds before withdrawing the needle from the skin
5. Lightly press gauze or cotton ball on the injection site as needed. Do not put the needle cap back on
6. GIVLAARI vials are for single use only. Discard unused portion of the drug. Dispose of needles, syringes, and needle caps in a locally approved sharps disposal container

In case of a missed dose¹

If a dose is missed, administer GIVLAARI as soon as possible.

Resume dosing at monthly intervals following administration of the missed dose.

IMPORTANT SAFETY INFORMATION

Injection Site Reactions

Injection site reactions were reported in 25% of patients receiving GIVLAARI in the placebo-controlled trial. Symptoms included erythema, pain, pruritus, rash, discoloration, or swelling around the injection site. One (2%) patient experienced a single, transient, recall reaction of erythema at a prior injection site with a subsequent dose administration.

Please see **Important Safety Information on page 3** and full **Prescribing Information**.



injection for subcutaneous use
189 mg/mL




How GIVLAARI® (givosiran) is supplied¹

- GIVLAARI is a ready-to-use solution supplied in single-dose vials of 189 mg/mL in cartons containing one vial



Storage and handling¹



Keep in carton



Store at 2°C to 25°C
(36°F to 77°F)



Do not freeze²

IMPORTANT SAFETY INFORMATION

Renal Toxicity

Increases in serum creatinine levels and decreases in estimated glomerular filtration rate (eGFR) have been reported during treatment with GIVLAARI. In the placebo-controlled study, 15% of patients receiving GIVLAARI experienced a renally-related adverse reaction. The median increase in creatinine at Month 3 was 0.07 mg/dL. Monitor renal function during treatment with GIVLAARI as clinically indicated.

Please see Important Safety Information on page 3 and full Prescribing Information.

 **GIVLAARI®**
(givosiran) injection for subcutaneous use
189 mg/mL



Provider Readiness Guide

Please see Important Safety Information on page 3 and full Prescribing Information.



Alnylam Assist™ is here to help

Alnylam Assist™ is dedicated to helping guide your patient through treatment with an Alnylam product.

Alnylam Assist™ offers support to help with:

- Securing access to an Alnylam product for your patient
- Initiating treatment for your patient
- Ordering product

For more information about how Alnylam Assist™ can help your patients, visit www.AlnylamAssist.com.

Preparing for the coverage and reimbursement process

When prescribing an Alnylam product, please refer to the steps below.

With payers

- 1 | Contact the payers through whom your patient has insurance coverage (commercial, local Medicare Administrative Contractor, State Medicaid, etc.) for additional information regarding appropriate coverage, coding, and payment policies.
 - For example, discuss the payment methodology for the appropriate Healthcare Common Procedure Coding System (HCPCS) code with payers and what constitutes a clean claim
- 2 |
 - Review the payer-specific coverage requirements and key medical necessity criteria
- 3 |
 - Ensure accurate and proper chart documentation

With your practice

- 4 | Know who in your practice is responsible for each of the following tasks:
 - Receiving benefit verification information
 - Submitting prior authorization/precertification, if required
 - Discussing financial obligations with patients
 - Scheduling appointments for drug administration
 - Ordering product for your patients
 - Submitting claims to payers

The Alnylam Assist™ team includes **Field Reimbursement Directors** who are knowledgeable in chart documentation best practices and billing and coding requirements for Alnylam products. They can answer your questions on these topics.

Please see **Important Safety Information** on page 3 and full **Prescribing Information**.





- 5 Update charge master/electronic billing system to ensure that the Alnylam product is recognized.
- 6 Anticipate requests from payers for clinical documentation if filing a claim for an Alnylam product.

Initiating therapy

When preparing to treat a patient with an Alnylam product at your practice, follow the steps below to help enable patient access, proper claims submission, and reimbursement.

- 1 Together with your patient, complete the Alnylam Assist™ **Start Form** to enroll your patient in Alnylam Assist™.
 - Your patient's Alnylam Case Manager will initiate verification of benefits and eligibility assessment for patient financial assistance, if appropriate.
 - To access the Alnylam Assist™ **Start Form**, visit www.AlnylamAssist.com
- 2 Schedule the patient for treatment.
- 3 Work with Alnylam Assist™ to determine the method for ordering product.
 - Alnylam Assist™ will send your patient's prescription to a specialty pharmacy and/or provide you with details about a specialty distributor. For some patients, home administration may also be an option depending on their insurance coverage.
- 4 After treatment, complete and submit the claim to the payer, if appropriate.



To get started, go to www.AlnylamAssist.com and complete the Alnylam Assist™ **Start Form** with your patient.

Please see Important Safety Information on page 3 and full Prescribing Information.

 **GIVLAARI®**
(givosiran) injection for subcutaneous use
189 mg/mL



Monday–Friday, 8AM–6PM ET

☎: 1-833-256-2748 | 📄: 1-833-256-2747

To learn more about GIVLAARI® (givosiran),
visit www.GIVLAARIHCP.com.

References: 1. GIVLAARI [package insert]. Cambridge, MA: Alnylam Pharmaceuticals, Inc; 2019. 2. Data on File. Alnylam Pharmaceuticals, Inc; 2019.

Please see **Important Safety Information** on page 3
and full **Prescribing Information**.



GIVLAARI, Alnylam Assist, and their associated logos are trademarks of Alnylam Pharmaceuticals, Inc.
© 2022 Alnylam Pharmaceuticals, Inc.

GIVLAARI®
(givosiran) injection for subcutaneous use
189 mg/mL

AS1-USA-00561-V2